



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A NEWLY OBSERVED STATION FOR *GALINSOGA*
HISPIDA.

SOME time ago the writer² called attention to the fact that besides *Galinsoga parviflora* Cav. and its var. *hispida* DC., another quite distinct species, *G. hispida* Benth., had been found in the Atlantic states, occurring at Camden, New Jersey, where it was collected on waste lands by Mr. C. F. Parker. This species was again found last October by Mr. Howard Schriver at several points in Cumberland, Maryland. Mr. Schriver, noticing its differences from *G. parviflora* Cav., sent specimens to the U. S. National Museum, whence they were kindly forwarded to the writer by Dr. J. N. Rose.

Mr. Schriver reports the plant as not only forming a large mass of vegetation (5 or 6 meters long) near a "bonded warehouse" or sort of cattle depot, but also extending down the banks of an adjacent stream where individuals were scattered on steep cliffy slopes of Helderberg limestone, even reaching the water's edge. A sporadic specimen was also found on a country road not far away. While the presence in the neighborhood of the cattle yard, a railway, and a distillery would readily account for the introduction of plants from a distance, the present species in the luxuriance of its growth and tendency to spread into the indigenous vegetation would suggest that it has found congenial conditions and is likely to persist. The species is readily distinguished by its "pink" or "red" rays (drying purple) and by its short pappus, which is about half the length of the achene. In foliage and rather copious spreading pubescence it resembles *G. parviflora*, var. *hispida* DC., but both *G. parviflora* and its variety have white rays and much longer pappus (nearly or quite the length of the achene).

As *G. hispida* Benth. is likely to be found at other stations in the middle Atlantic states it may be worth while to cite its more important synonymy, which is as follows:

Vargasia caracasana DC. Prodr. 5: 676. 1836.

Galinsoga hispida Benth. Bot. Sulph. 119. 1844.

G. brachystephana Regel, Ind. Sem. Hort. Turic. 2. 1846.

G. caracasana Sch. Bip. Linnaea 34: 529; also Bull. Soc. Bot. Fr. 18: 86. 1865.—B. L. ROBINSON, *Gray Herbarium*.

² Proc. Am. Acad. 29: 326.